

Patient Name:

Chart ID:

Address:

Phone Number:

DOB:

*Indications – Prescription treatment for hypotrichosis used to grow eyelashes.*

LATISSE SCREENING INTERVIEW

- |  |                              |                             |
|--|------------------------------|-----------------------------|
| 1. Do you use Lumigan or any other medication to treat intraocular pressure? | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| 2. Are you pregnant or breastfeeding?  | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| 3. Do you experience any of the following symptoms / conditions:             |                              |                             |
| a. Eye pruritus (itchy eyes)   | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| b. Conjunctival hyperemia (red eyes)   | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| c. Irritated eyes  | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| d. Dry eyes  | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| e. Eyelid redness  | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| f. Twitching eyelids   | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| g. Macular edema   | <input type="checkbox"/> Yes | <input type="checkbox"/> No |
| 4. Do you have brown spots or discoloration of the eyelids?                  | <input type="checkbox"/> Yes | <input type="checkbox"/> No |

*If you answered Yes to #1 or #2, you may not use Latisse.*

*If you answered Yes to any part of #3, we recommend you see your Ophthalmologist to correct and control your symptoms prior to starting Latisse.*

*If you answered Yes to #4, we recommend you see your Dermatologist before starting Latisse.*

Latisse warnings, precautions and adverse reactions:

1. Do not use Latisse with other medications that lower Intraocular Pressure (IOP).
2. May cause permanent iris pigmentation if Latisse comes in continual contact with your eye.
3. Latisse may cause hyper pigmentation of the eyelid. This is reversible after stopping Latisse.
4. It is very important to apply Latisse to skin of upper eyelid margin at the base of the eyelashes to reduce the potential of unwanted hair growth outside treatment area.
5. Do not use Latisse if you have Macular edema.
6. Avoid contaminating Latisse applicators.
7. Do Not use if pregnant or breastfeeding.
8. Remove contact lenses prior to applying Latisse. Resume use of contact lenses 15 minutes after Latisse dries.
9. Red eye or conjunctal hyperemia will usually resolve in a few days to two (2) weeks. If it persists or suddenly worsens, discontinue Latisse.
10. Less than 4% of patients experience eye pruritus (itch), skin hyper pigmentation (darkening), ocular (eye) irritation, dry eye, Erythema (redness) of eyelids.

*If these symptoms do not resolve in a few days, then discontinue Latisse and / or see your Ophthalmologist*

\*\*\*Non-refundable – Latisse is a prescription drug – cannot be returned\*\*\*

Patient Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Witness Signature: \_\_\_\_\_

Date: \_\_\_\_\_

**Use as directed – see package insert.**